

9600157

## THE CONTRED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

# NASH Research Joundation

Thereis, there has been presented to the

#### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED, PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT-TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR PRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR ING IT FOR ANY OF THE ABOVE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. IN STATES SEED OF THIS VARIETY (I) SHALL BE SOLD BY VARIETY NAME ONLY AS A CLASS OF AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF

WHEAT

'Ernest'

In Testimone Marcost, I have hereunto set my hand and caused the seal of the Hant Variety Protection Office to be affixed at the City of Washington, D.C. this turnty-righth day have in the year of our Lord one thousand

Allest.

Marsha d. Fun ton

Commissioner
Plant Variety Protection Office
Aurica Aurica Washeling Service


Genetary of Syriculture

REPRODUCE LOCALLY. Include form number and date	e on all reproduction	<b>15</b> .		FORM APPROVE	ED - OMB NO. 0581-005
U.S. DEPARTMENT OF AGRICULTUM AGRICULTURAL MARKETING SERVI SCIENCE DIVISION - PLANT VARIETY PROTEC	The following statements are made in accordance with the Privacy Act 1974 (5 U.S.C. 552a).  Application is required in order to determine if a plant variety protect certificate is to be issued (7 U.S.C. 2421). Information is held confident until certificate is issued (7 U.S.C. 2426).				
APPLICATION FOR PLANT VARIETY PROTE					
1. NAME OF APPLICANT(S) (as it is to appear on the Certificate)		-	2. TEMPORARY DESIGNATION OR	3. VARIETY NA	ME
NDSU Research Foundation	•		EXPERIMENTAL NUMBER ND6.77	Ernest	
NDSO Research Foundation			ND0,77	Linest	
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, 4	and Country)		6. TELEPHONE (include area code)		
c/o Executive Director PO Box 5014 Fargo, ND 58105-5014	ana country	7	701-231-8931	PAGES AND THE PROPERTY.	9600157
14190, 112 00100 0011	•		6. FAX (include area code)	F DATE	
			701-231-1013	2 2-1	396
7. GENUS AND SPECIES NAME	8. FAMILY NA	ME (Botanio	al)	FILING AND	EXAMINATION FEE
<u>Triticum</u> <u>aestivum</u> L.	Grami	neae		F - 24	0.00
9. CROP KIND NAME (Common name)				DATE 2	0.
Hard red spring wheat				2 3-16	F76
10. IF THE APPLICANT NAMED IS NOT A "PERSON", GIVE FORM OF OR 501 (c) (3) Corporation - NDSU R				C GERTIFICATI	Office Control of the
11. IF INCORPORATED, GIVE STATE OF INCORPORATION	escaron roun		12. DATE OF INCORPORATION	V E DATE	
North Dakota				0	9.47
13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY,				14. TELEPHONE	(include area code)
Richard C. Froberg Department of Plant Sciences		e Zeto	cna Director	701-231	-8143
North Dakota State University			arch Foundation	15. FAX (include	area codel
PO Box 5051	P0	Box 50	14	1	· · ·
Fargo, ND 58105-5051			58105-5014	701-231	-04/4
<ol> <li>CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (</li> <li>Exhibit A. Origin and Breeding History of the Veriety</li> </ol>	Follow instructions on reve	erse)	-		
b. X Exhibit B. Statement of Distinctness					
c. X Exhibit C. Objective Description of the Variety					
d. 🔀 Exhibit D. Additional Description of the Variety  e. 🔀 Exhibit E. Statement of the Basis of the Applicant's Ownershi	in.				
f. Noucher Sample (2,500 viable untreated seeds or, for tuber pa		tion that tiss	ue culture will be deposited and maintain	ed in a public reposit	toryl
g. 🛛 Filing and Examination Fee (\$2,460), made payable to "Treast	rer of the United States" /	Mail to PVP	0)		,
7. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE SO  X YES (If "yes," answer items 18 and 19 below)		NLY, AS A		n 83(a) of the Plant	Variety Protection Acti?
8. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE LI GENERATIONS?	IMITED AS TO NUMBER O	F 19.	F "YES" TO ITEM 18, WHICH CLASSES	OF PRODUCTION B	EYOND BREEDER SEED?
Ø YES □ NO			S FOUNDATION REGISTERS		· <del>-</del>
<ul> <li>HAS THE VARIETY OR A HYBRID PRODUCED FROM THE VARIETY B.</li> <li>YES III "yes," give names of countries and dates!</li> </ul>	□ NO USA	- Rel	sale, on Marketed in the U.S. on o ease date March 10, st seed sale Novemb	1995	
<ol> <li>The applicant(s) declare that a viable sample of basic seed of the variety applicable, or for a tuber propagated variety a tissue culture will be dependent.</li> </ol>	y will be furnished with apposited in a public reposited	plication and	will be replenished upon request in according for the direction of the confidence	dance with such req	gulations as may be
The undersigned applicant(s) is(are) the owner(s) of this sexually reprod Section 41, and is entitled to protection under the provisions of Section	uced or tuber propagated r	alant variety	and helievelet that the variety is near at	stinct, uniform, and	stable as required in
Applicant(s) is(are) informed that false representation herein can jeopard	•				
SONATURE OF APPLICANT (Owner(s))		SIGNATURE	OF APPLICANT (Owner(s))		
AME (Please print or tylph)		NAME IPIA	se print or type)		
Dale Zetocha					
Executive Director NDSU Research Foundation	3/12/96	CAPACITY	OR TITLE		DATE
ABOU NESCUTON FOUNDATION					<u> </u>

## EXHIBIT A - ORIGIN AND BREEDING HISTORY 'ERNEST'

Spring 1985	Original cross made at North Dakota State University (NDSU) greenhouse. Cross number is 85373 Pedigree - ND622*2/Cutless ND 622 = SU7 sib-3//SU28-1*3/Agent
Summer 1985	F <sub>1</sub> plants, NDSU greenhouse.
Summer 1986	${\bf F_2}$ bulked population, NDSU research land.
Summer 1987	${\tt F_3}$ bulked population, NDSU research land.
Spring 1988	${\sf F_4}$ single-seed-descent population, NDSU greenhouse.
Summer 1988	$F_5$ plant row, NDSU research land.
Summer 1989	$F_6$ progeny plot, NDSU research land.
Summer 1990	$F_7$ preliminary yield trial, 3 locations, NDSU research land.
Summer 1991	F <sub>8</sub> elite yield trial, 4 locations, NDSU research land. Seed increase (37 lbs.) Prosper, ND.
Summer 1992-1994	North Dakota HRS wheat variety trial, 4 locations each year. North Dakota HRS wheat off-station trial, 4 locations each year.
	Experimental line designation - ND677
Summer 1992-1994	Seed increase by NDSU Seedstocks Project.
Summer 1993-1994	Uniform Regional HRS Wheat Nursery, about 20 locations each year, Upper Midwest/U.S.A.
Summer 1993-1994 March 10, 1995 March 17, 1995	Wheat Quality Council test Release date of Ernest. Seed of 'Ernest' (experimental line ND677) distributed to contract seed growers.

Observations indicate Ernest is uniform and stable within commercially acceptable limits for all traits as described in Exhibit C. Uniformity and stability of Ernest were observed 1989 to 1994 for six generations ( $F_6$  -  $F_{11}$ ). Variants (taller plants, 5-12 cm) occurrence at a frequency of 5/10,000 and awnless plant variants at a frequency of 2/10,000.



North Dakota State University
Loftsgard Hall
P.O. Box 5051
Fargo, North Dakota April 11, 1996
58105-5051 USA

Tel.701.231.7971 Fax701.231.8474

> Dr. Alan A. Atchley Plant Variety Protection Office NAL Building, Room 500 10301 Baltimore Blvd. Beltsville, MD 20705

Biotechnology Breeding Forestry Genetics Horticulture Physiology Production Weed Science

SUBJECT: PVP Application No. 9600157, Wheat, 'Ernest'

Dear Dr. Atchley:

Following is the information that you requested (letter 3 April 1996) so that the examination of 'Ernest' wheat may be concluded:

The Variety Release Committee of the North Dakota Agricultural Experiment Station (NDAES) recommended to Dr. Robert Todd, Director, NDAES, that the experimental wheat line ND677 be named Ernest and released as a cultivar. Dr. Todd announced this release through public media on March 10, 1995. Seed of Ernest was distributed by the NDAES through the North Dakota Crop Improvement county groups to experienced seed producers for seed multiplication during the growing season of 1995. These producers were under contract to the Seedstocks Project, Department of Plant Sciences, NDAES, North Dakota State University for the Ernest seed produced. On November 15, 1995, these contracted seed producers were released from their contract and Ernest seed was then available and offered for sale to the public in the U.S.A.

#### Exhibit A

'Cutless' is a commercial cultivar developed and released by the North Dakota Agricultural Experiment Station, North Dakota State University. SU7 and SU28 are public lines developed by Dr. Glenn S. Smith as an employee of North Dakota State University.

Selection criteria for the breeding of Ernest wheat were highly heritable traits (i.e., plant height, maturity, disease resistance) in the early segregating



Dr. Alan A. Atchley Page 2 April 11, 1996

generations,  $F_2$  -  $F_5$ . Beginning in the  $F_6$  generation, selection criteria also included lodging resistance, shattering resistance, reaction to wheat stem sawfly, test weight, grain yield and bread making characters (grain protein, milling extraction, dough mixing, loaf volume, etc.). Data were obtained from multiple locations and over years to evaluate and identify the experimental line ND677 that was named Ernest. Overall, the selection criteria were a combination of traits used to identify a superior hard red spring wheat genotype adapted to northwestern North Dakota wheat production where the wheat stem sawfly is a frequent production hazard, and a genotype having acceptable (compared to check cultivars) milling and bread making properties for domestic and export markets

and export markets.

[Exhibit Al: ] AAA laMay 1996

Difference between Ernest and Butte 86

Ernest has solid stem internodes. Butte 86 has hollow stem internodes.

This information should provide facts for the examination of Ernest wheat.

Sincerely,

Richard C. Frohberg

**Professor** 

RCF:lh

#### EXHIBIT B - NOVELTY STATEMENT

To my knowledge 'Ernest' most nearly resembles 'Amidon'. Differences include, but are not necessarily restricted to, the following:

1. Ernest has solid stem internodes. Amidon has hollow stem internodes.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE COMMODITIES SCIENTIFIC SUPPORT DIVISION BELTSVILLE, MARYLAND 20705

EXHIBIT C

## OBJECTIVE DESCRIPTION OF VARIETY

	RITICUM SPP.)
NDSU Research Foundation ADDRESS (Street and No. or R.P.D. No., City, State, and ZIP Code)	POR OFFICIAL USE ONLY-7
PO Box 5014	VARIETY NAME OR TEMPORARY DESIGNATION
Fargo, ND. 58105 - 5014	, and the second
Place the appropriate number that describes the varietal charact Place a zero in first box (e.s. 0 8 9 or 0 9 ) when number	er of this variety in the boxes below. is either 99 or less or 9 or less.
I. KIND:	
	5 = POLISH 6 = POULARD 7 = CLUB
2. TYPE,  1 1 = SPRING 2 = WINTER 3 = OTHER (Specify)	1 = SOFT 3 = OTHER (Specify) 2 = HARO
2 1 = WHITE 2 = RED 3 = OTHER (Specify)	<del>-</del>
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	
057 FIRST FLOWERING	0 6 0 LAST FLOWERING
4. MATURITY (50% Flowering):	
03 NO. OF DAYS EARLIER THAN	. 3 l = ARTHUR 2 = SCOUT 3 = CHRIS
NO. OF DAYS LATER THAN	A = LEMHI 5 = NUGAINES 6 = LEEDS
S. PLANT HEIGHT (From soil level to top of head):	
0 9.4 cm. HIGH	
CM. TALLER THAN	
0 7 CM. SHORTER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS  4 = LEMNI 5 = NUGAINES 6 = LEEDS
6. PLANT COLOR AT BOOTING (See reverse):	7. ANTHER COLOR:
3 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN	1 = YELLOW 2 = PURPLE
8. STEM	
Anthocyanin: 1 = ABSENT 2 = PRESENT	Waxy bloom: 1 = ABSENT 2 = PRESENT
Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT	2 Internodes: 1 = HOLLOW 2 = SOLID
0 4 NO. OF NODES (Originating from node above ground)	26 CM INTERNODE LENGTH BETWEEN FLAG LEAF AND LEAF BELOW
AURICLES:	
Anthocyanin: 1 = ABSENT 2 = PRESENT	Hairiness: 1 = ABSENT 2 = PRESENT
0. LEAF:	
Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):	2 Flag leaf: 1 = NOT TWISTED 2 = TWISTED
Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT	2 Waxy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT
0 9 MM. LEAF WIDTH (First leaf below flag leaf)	23 CM. LEAF LENGTH (First leaf below flag leaf):
ORM LMGS 470-6 (6-82) (Formerly Form   PGS 470-6 (3-76) which m	who wood

Density: 1 = LAX	2 = DENSE	Shape: 1 = TAPERI 4 = OTHER	NG 2 = STRAP 3 = CLAVATE (Specify)
4 Awnedness: 1 = AWN	LESS 2 = APICALLY AWNLETED 3	= AWNLETED 4 = AWNE	• .
2 Color at maturity: 5 =	WHITE 2 = YELLOW 3 = PINK 4 = BROWN 6 = BLACK 7 = OTHE	: REO R (Specity):	•
06 CM. LENGTH		MM. WIDTH	•
12. GLUMES AT MATURIT  Length: 1 = SHORT (  3 = LONG (	CA. 7 mm.) 2 = MEDIUM (CA. 8 mm.)	2 Width: 1 = NARROW 3 = WIDE (C.	
Shoulder 1 = WANTII shape: 4 = SQUAR		3 Beak: 1=08TUSE	2 = ACUTE 3 = ACUMINATE
13. COLEOPTILE COLOR:		14. SEEDLING ANTHOCY	AHIH:
1 = WHITE 2 = RE	D 3 = PURPLE	1 = ABSENT 2	= PRESENT
15. JUVENILE PLANT GRO	OWTH HABIT:		
2 1 = PROSTRATE	2 = SEMI-ERECT 3 = EREC	т	
16. SEED:			
2 Shape: 1 = OVATE	2 = OVAL 3 = ELLIPTICAL	2 Cheek: 1 = ROUND	ED 2 = ANGULAR
Brush. 1 = SHORT	2 = MEDIUM 3 = LONG	Brush: 1 = NOT CO	DLLARED 2 = COLLARED
Phenol reaction (See instructions):	1 = IVORY 2 = FAWN 3 = LT. BROWN 4 = BROWN 5 = BLACK	•	· ·
3 Color: 1 = WHITE	2 = AMBER 3 = RED 4 = PURPLE	5 = OTHER (Specity)	
D 6 MM. LENGTH	△ 3 MM. WIDTH	3 2 GM. PER 1000	SEEOS
IT. SEED CREASE:			
12	ESS OF KERNEL 'WINOKA'	1.2   "	LESS OF KERNEL 'SCOUT'
2 = 80% OR LE	SS OF KERNEL 'CHRIS'		LESS OF KERNEL "CHRIS"
	S WIDE AS KERNEL 'LEMHI'	J = 30 % O A	LESS OF REINIGE. LEWIS
2 STEM RUST QFBS, GRACOO, RTQQ,	AC TEST CONTROL (Recent Local	STRIPE RUST	O LOOSE SMUT
D POWDERY MILDEW	O BUNT	OTHER (Specify)	
19. INSECT: (0 = Not Teste	d, 1 = Susceptible, 2 = Resistant)		
2 SAWFLY	O APHID (Bydv.)	Ø GREEN BUG	O CEREAL LEAF BEETLE
OTHER (Specily)	HESSIAN FLY	0 GP 0 A	0 s 0 c
•	RACES:	0 o 0 E	0 , 0 ,
20. INDICATE WHICH VARIE	TY MOST CLOSELY RESEMBLES THAT S	UBMITTED:	
CHARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Plant tillering	Amidon	Seed size	Amidon
Leaf size	Amidon	Seed shope	Amidon
Leaf color	Amidon	Coleoptile elongation	Amidon
Leaf carriage	Amidon	Seedling pigmentation	Amidon

#### INSTRUCTIONS

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

## EXHIBIT D - ADDITIONAL DESCRIPTION OF VARIETY

When 'Ernest' is compared to 'Stoa', 'Amidon' and 'Grandin' it has wheat stem sawfly resistance as determined by stem cutting percentage and is similar to 'Lew' and 'Cutless' (Table 10).

### ERNEST HRS WHEAT

Table 1. Mean grain yield of selected entries in the 1992-94 hard red spring wheat variety trial at North Dakota Agricultural Research Centers.

Variety or line	Dickinson†	Hettinger‡	Williston	North Central
		bu/A	_	
Conv. ht.				
Stoa	46.6	68.9	54.3	47.9
Butte 86	41.1	70.5	52.1	45.6
Amidon	47.4	74.0	55.9	44.9
Sharp		70.1	53.0	44.6
CDC Teal	42.1	65.8	53.3	45.8
AC Domain	35.7		49.3	41.9
Kulm		74.5	55.4	45.3
Fortuna		-	44.3	·
Lew	·		51.3	
Leader	35.8	40 pm - 40 mg	48.2	
Ernest	50.8	70.1	53.7	42.7
<u>Semidwarf</u>			·	
Cutless		41.7	46.6	
Rambo	40.2		53.4	
Glenman			53.6	
2375	45.1	69.8	57.6	46.1
Grandin	46.8	68.8	56.5	47.1
Norm		77.4	56.9	47.5
Hi Line		65.0	55.1	39.6
McNeal	55.1	88.6	63.1	49.9

<sup>†1993</sup> yield data

<sup>‡1992</sup> and 1993 yield data

Table 2. Summary of agronomic performance of selected entries in the 1992-94 hard red spring wheat variety trial at North Dakota Agricultural Research Centers†.

Variety or line	Headed (10)‡	Height (10)	Lodging score (1)	Leaf disease (8)	Test weight (10)	Grain yield (8)
	days	cm	0-9	%	lb/bu	bu/A
Conv. Ht.						
Stoa	65	88	0.3	40	60.0	56.0
Amidon	65	87	0.3	30	60.5	56.5
CDC Teal	63	85	1.8	36	60.2	53.5
Kulm	61	83	0.3	50	62.1	56.5
Ernest	65	86	0.3	31	61.6	53.9
Semidwarf						
2375	62	79	2.2	50	60.8	56.9
Grandin	64	81	0	45	61.2	56.4
Norm	63	77	0.5	26	60.0	58.9
McNeal	66	80	0.8	28	60.2	64.3

<sup>†</sup>Dickinson, Hettinger, Williston, North Central.

<sup>‡</sup>Number of tests each trait.

Table 3. Grain yield of selected entries in 1992-94 hard red spring wheat trials conducted by the Williston Research Center.

		Wil	lliston		Off-station locations†		
Variety or line	Variety	Early fallow	Recrop	Notill	1992 <sup>-</sup> (5)	1993 (4)	1994 (4)
				— bu/A —			
Conv. Ht.							•
Stoa	54.3	56.9	44.1	41.5	57.9	55.4	42.6
Butte 86	52.1				58.5	52.6	40.9
Amidon	55.9	59.5	48.6	44.6	64.2	54.1	43.5
CDC Teal	53.3				<b></b>		39.5
Kulm	55.4				55.9		44.5
Ernest	53.7	54.7	46.5	44.0	60.3	58.0	42.6
Semidwarf							
2375	57.6	58.3	49.3	45.0	64.4	55.5	47.7
Grandin	56.5	56.3	45.5	42.3	60.6	52.5	42.4
Bergen	61.8	59.8	51.4	49.1	68.1	61.3	47.6
Dalen	56.9	56.8	49.7	44.8	66.5	56.7	43.9
Norm	56.9				64.9	54.4	
Rambo	53.5					44.9	35.2
Hi Line	55.1					48.2	40.9
McNeal	63.1				er m m =	60.6	46.0

†Locations each year:

1992 - Powers Lake, Arnegard, Newtown, Stanley, Fortuna.

1993 - Arnegard, Newtown, Stanley, Fortuna.

1994 - Arnegard, Newtown, Stanley, Burke County.

Table 4. Summary of agronomic performance of selected entries in the 1992-94 hard red spring wheat trials conducted by the Williston Research Center.

Variety or line	Headed (8)†	Height (17)	Lodging score (2)	Leaf disease (11)	Test weight (26)	Grain yield (25)
	days	cm	%	%	lb/bu	bu/A
Conv. Ht.	÷					÷
Stoa	62	93	2	43	59.2	51.0
Amidon	62	91	2	29	59.6	53.4
Ernest	61	91	2	33	60.9	52.0
Semidwarf						
2375	60	86	28	52	60.6	54.7
Grandin	61	83	0	48	59.8	51.4
Bergen	61	77	0	18	60.1	57.7
Dalen	59	75	6	34	60.9	54.4

<sup>†</sup>Number of tests each trait.

Table 5. Grain yield of selected entries in 1993-94 hard red spring wheat trials conducted by the North Central Research Center.

		•			locations†
Variety		Minot		1993	1994
or line	Fallow	Recrop‡	Notill	(2)	(4)
			— bu/A ———		
Conv. Ht.					
Stoa	54.2	43.1	52.8	41.1	48.1
Butte 86	52.2	41.1	52.7	39.7	40.8
Amidon	51.9	42.4	52.4	36.6	37.8
Kulm	52.1	37.2	49.1	43.2	44.5
				-	
Ernest	50.9	41.4	52.2	41.2	43.4
<u>Semidwarf</u>					
2375	53.0	37.6	52.5	39.9	46.9
Prospect	56.6	35.8	53.8	47.7	50.9
Gus	54.4	35.8	50.2	34.6	40.4
Grandin	55.2	42.7	53.2	36.8	44.7
2371	53.5	37.7	52.5	41.4	42.1
Dalen	55.9	36.0	53.5	44.0	42.2
Norm	56.0	45.9	55.5	46.0	45.0

†Locations each year:

1993 - Harvey, York 1994 - Fessenden, Maddock.

‡1993 data.

Table 6. Summary of agronomic performance of selected entries in the 1993-94 hard red spring wheat trials conducted by the North Central Research Center.

Variety or line	Headed (6)†	Height (10)	Lodging score (4)	Leaf disease (5)	Test weight (10)	Grain yield (9)
	days	cm	0-9	%	lb/bu	bu/A
Conv. Ht.	•					
Stoa	67	103	2.0	42	58.5	49.2
Butte 86	63	93	0.7	62	59.4	46.2
Amidon	66	98	3.3	32	58.6	44.8
Kulm	62	97	0.7	69	60.4	46.5
Ernest	66	99	1.6	37	59.7	46.8
<u>Semidwarf</u>						
2375	64	89	3.9	60	59.1	47.7
Prospect	66	90	0.4	53	58.3	51.1
Gus	67	93	1.8	34	58.0	44.6
Grandin	64	89	0.4	64	59.6	47.6
2371	65	82	0.9	53	58.1	46.8
Dalen	64	80	0.6	53	59.5	48.0
Norm	65	86	0.4	36	58.1	50.5

<sup>†</sup>Number of tests each trait.

Table 7. Summary of agronomic performance of selected entries in 1994 hard red spring wheat trials conducted by the Hettinger Research Center.

	-		Location - grain yield					
Variety or line	Height (4)	Test weight (4)	Scranton	Selfridge	Regent	New Leipzig	x (4)	
	cm	lb/bu			bu/A			
Conv. Ht.								
Amidon	76	60.2	44.0	48.1	42.8	36.7	42.9	
Kulm	73	61.7	37.3	56.3	47.9	36.9	44.6	
Ernest	72	60.4	45.5	52.6	46.3	33.1	44.4	
Semidwarf				· ·				
2375	69	60.3	50.6	59.2	50.5	36.8	49.3	
Gus	67	59.5	47.8	50.0	49.9	35.8	45.9	
Grandin	67	60.4	43.0	44.6	42.2	33.1	40.7	
2371	63	58.7	49.4	57.0	49.0	35.3	47.7	
Dalen	61	61.4	47.7	46.5	48.0	39.4	45.4	
Krona	62	59.4	55.9	64.1	48.7	47.3	54.0	
Norm	63	60.3	49.2	67.1	50.4	40.2	51.7	
Sonja	62	59.4	44.4	64.7	56.0	39.6	51.2	
McNeal	70	59.8	46.6	56.9	52.6	40.5	49.2	
C.V.%			10.6	7.1	9.5	6.7	ē.	
LSD (.05)			7.4	5.7	6.7	4.3		

Table 8. Summary of agronomic performance of selected entires in the 1993 hard red spring trials conducted by the Williston Research Center (Lew vs. Ernest comparison).

Variety or line	Headed (4)†	Height (4)	Lodging (2)	Leaf disease (4)	Test weight (4)	Grain yield (4)
	days	cm	%	%	lb/bu	bu/A
Conv. Ht.						
Stoa	61	84	2	47	57.6	56.3
Amidon	61	78	2	33	57.7	59.1
Lew	63	84	66	45	59.1	48.4
Ernest	60	78	2	29	58.5	55.4
Semidwarf		·				
2375	59	75	28	58	58.0	60.2
Grandin	60	76	0	53	58.4	58.1
Norm	61	74	3	19	57.3	59.9
Rambo	63	72	1	45	58.1	51.4

<sup>†</sup>Number of tests each trait.

Table 9. Summary of agronomic performance of selected entries in the 1993-94 hard red spring wheat variety trial at North Dakota Agricultural Research Centers† (AC Eatonia vs. Ernest comparison).

Variety or line	Headed (6)‡	Height (6)	Lodging score (1)	Leaf disease (5)	Test weight (6)	Grain yield (5)
•	days	cm	0-9	%	lb/bu	bu/A
Conv. Ht.					·	•
Stoa	64	92	0.3	51	59.1	57.7
Amidon	65	90	0.3	40	60.1	57.5
CDC Teal	64	89	1.8	39	59.4	52.8
AC Eatonia	65	90	4.3	51	60.0	47.9
Ernest	64	90	0.3	41	61.1	56.4
Semidwarf						
Grandin	63	83	0	59	60.1	58.9
Norm	64	80	0.5	34	59.2	61.8
McNeal	66	83	0.8	37	59.5	64.6

<sup>†</sup>Dickinson, Hettinger, Williston, North Central

<sup>‡</sup>Number of tests each trait.

Table 10. Stem cutting percentage (wheat stem sawfly) of selected entries in hard red spring wheat trials conducted by the Williston Research Center, 1991 and 1992.

	Williston say	wfly nursery <sup>‡</sup>
Variety or Line	1991	1992
	%	%
Stoa	88	76
Amidon	63	66
Grandin	90	68
Lew	13	13
Cutless	7	18
Leader		24
Ernest	17	26
LSD.05	24	14
CV%	18	22

<sup>&</sup>lt;sup>†</sup> No stem cutting in 1993 and 1994 trials. <sup>‡</sup> Visual estimates - 1991; actual stems cut by sawfly (%) - 1992

Table 11. Summary of agronomic performance of selected entries in the Uniform Regional Hard Red Spring Wheat Nursery, 1993 and 1994.

1993

	4	4	Lodging	R	Rust <sup>†</sup>		Test	Grain
Variety or line	Headed (15) <sup>‡</sup>	Height (16)	Score (7)	Leaf (1)	Stem (1)	Shattering (1)	weight (16)	yield (17)
•	days	cm	1-9	%	%	%	lb/bu	bu/A
Stoa	34	100	2.9	10MR-MS	0	15	56.0	50.1
Butte 86	30	92	2.8	10MR-MS	5R-MR	5	57.4	52.8
Kulm	30	96	2.0	tR-MR	tR	15	58.3	52.7
Trenton	32	103	2.7	20MR	30MR-MS	5	57.3	49.4
Ernest	32	96	2.6	tR	0	5	56.9	46.2
F-test	52.6	70.0	4.3				13.3	8.7
LSD.05	0.9	3.0	0.9	***		·	1.2	5.5
CV%	3.7	4.8	30.4			***	3.2	16.3

<sup>†</sup> Data from field inoculated rust nursery, USDA-ARS, Cereal Rust Laboratory, St. Paul, MN.

<sup>‡</sup> Number of tests each trait

1994

			Lodging	R	lust <sup>†</sup>	Leaf	Test	Grain
Variety or line	Headed (18) <sup>‡</sup>	Height (19)	Score (7)	Leaf (1)	Stem (1)	Spotting (3)	weight (20)	yield (20)
	days	cm	1-9	%	%	%	lb/bu	bu/A
Stoa	27	92	3	<b>5</b> S	5R-MR	5	57.7	48.0
Butte 86	24	86	3	<b>5</b> S	10MR	8	59.4	47.8
Trenton	27	94	3	·	30MR-MS	9	59.0	47.2
Ernest	27	90	2	5MS-S	tR	22	59.0	42.6
F-test	49.9	99.9	3.4			1.8	12.6	16.1
LSD.05	0.8	2.0	1.2			24.4	0.8	2.9
CV%	4.4	3.9	47.3	***		95.3	2.2	9.9

<sup>†</sup>Data from 1994 rust nursery, USDA-ARS, Cereal Rust Laboratory, St. Paul, MN. ‡Number of tests each trait

NORTH DAKOTA STATE UNIVERSITY	AGRICULTURAL EXPERIMENT STATION	DEPARTMENT OF CEREAL SCIENCE AND FOOD TECHNOLOGY	ANALYTICAL, MILLING AND BAKING DATA	FIELD PLOT VARIETIES	(2 YEAR AVERAGE, 8 LOCATIONS)	1992 AND 1993
-------------------------------	---------------------------------	--	-------------------------------------	----------------------	-------------------------------	---------------

AVERAGES OF 2 IEARS	7	DADTHOODAY	•	!	:
		FARINOGRAM	~,	HOUR	3 HOUR FERMENTA

					210000											
	•	TEST	VIT	FAL	14% MB	FLR	WET	FLR	PEAK MIX		٠	MIX		LOAF		
VARIETY OR NUMBER AMIDON BUTTE86 GRANDIN LEN LEN STOA	YLD BPA ND ND ND ND	WT LB/BU 61.2 61.4 61.8 60.3 60.5	X 8888 X 888 X 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	NO SEC 412 1 420 1 407 1 388 1 419 1 1 419	WHT FLR 14.5 13.8 6 14.0 14.9 14.0 14.8 13.8 6 14.4 13.8 6 14.8 14.8 14.8 14.0 6	EXT 69.2 68.8 70.6 69.7 69.7	GLU % 42.7 43.3 40.2 39.0 41.3	EXT GLU ASH 69.2 42.7 0.38 68.8 43.3 0.39 70.6 40.2 0.40 69.7 39.0 0.40 69.9 41.3 0.38	TIME TOL N MIN MIN E 8.7 14.0 9.9 10.6 9.8 20.2 13.8 25.7 12.1 15.7	111 30 27 27 28 19 113 113	CLASS 5.7 5.4 6.9 6.7	ABS TIME % MIN 67.1 1.70 69.3 1.55 68.4 1.85 67.0 2.20 64.5 1.75	66 9.8 55 9.9 0 10.0	CC CC 851 841 839 893	 CB CT CL CL 8.2 10.0 8.2 10.0 7.7 10.0 8.5 10.0	T. SYH .0 10.0 .0 9.7 .0 9.8 .0 10.0

THE VALUES WERE DETERMINED BY AVERAGING VALUES FROM THE POLLOWING STATIONS: DI MI	R DETERNI ALUBS FRO STATIONS:	INED OM : DICKII MINOT WILLI	ED DICKINSON MINOT WILLISTON		NOI AGR DEP ARTHENT ANAL Y	NORT AGRIC THENT C	H DAI CLTUB P CR CAL, FIBLI	DAKOTA ST TURAL EXP CERBAL SC L, HILLIN ELD PLOT 1994 TABL	ATE UN ERIHEN IENCE G AND VARIET CROP E 13	II VERSIT TI STATI AND FOO BAKING TES	IVERSITY T STATION AND FOOD TECHNOLOGY BAKING DATA IES	)Logy						N. C.	
		•							,	PARINOGRAH	Ŧ		C	HOUR 1	<b>FERHENTATION</b>	TATIO	_		
	TEST	ST VIT	WHT T FAL		BIN MB	FLR	WET	FLR	PBAK HIX	×			HIX	1	LOAF			<u>[</u>	
VARIETY OR YLD		WT KER	R NO	H.	FLR	EXT (	פרה ז	ASH	TIME TOL	L MTI	61.199	ABS	TIHE	Q <b>X</b>	۔	85 E	៦៩	7	
AIN		9		15	9 6	67.6	44.8	7.0	~ u	0.0	7.3	63.8	in in	00	852 7		86	-	<u>.</u> .
DON	99			1	1	- lc	9	0.41	نان	.5 25	7.3	63.6	, 0	-	ما	ن ا	8 10.0	7	.1.
BERGEN ND	53	98 6.		13.4	12.6	70.1	36.6 0	0.43	17.7 19	.5 18	7.7	63.3	1.90 1	10.0	00	∞ •	.2 10.0	6.0	_
RLIN	9	.0 91	42	8	14:1		. 0	0.41	28.7 27	- <del></del>	7.7AB	62.7					7 10.0	10	_
SAL	59	.9		15.4	14.7	67.7	œ	0.40	28.8 32	.3 12	8.0AB	63.2		_	700	<b>40</b> (-	5 10	20	
*BRNEST ND	0 61.	96		14.3	13.4		150	0.38	.0 28	.0 13	7.3	59.9	101	? •	_	. ec	30.00	. 6	1
07	59	.0 88		14.0	13.4	66.7	10.5	5.43	15.7 22	.5 13	7.7	63.3	2.00	9.7	7	.0	0 10.0	8.3	١.
GRANDIN ND	09	.2 96	416	14.2	13.4	68.7	38.6	).42	26.2 29	.7 17	7.3AB	64.9	2.60 1	0.0	373 8	€	.0 10.0	9.0	
GUS ND	53	.7 98		15.0	14.2	67.3	41.7 0	0.42	13.3 22	.0 15	7.3	63.9	2.15	L. 6	903 8	0.0	2 10.0	م ه	_
-	9 40	6 92	407	<b>-</b>	?	0.69		. 7	•	.8 18	, .	59.5	2 10		9 40	- [-	101	י ס	
•	62	1		14.	6	67.4	9	<u> </u>		.2 13	•	64.7	10	0.	80	60	5 10	•	
co.	200	7 94		13		<u>.</u>	33.3	7.	30.5 24	 	7.0AB	59.4	2.	10.0	т с С с	<b>&amp;</b> F	2:	٠ د	
*HCNBAL ND	50	.5 96	404	13.5	12.6	64.9		0.40		.8.	• •	65.6	4.50 1		982 9	·	2010	-	_ 1
	09	ŀ		14.9	7.	9.	o .	7	.3 34	S.	8.0AB	64.7	in i	0.	957 8		10.	9.0	
678	62.	.a 98	331	14.2	13.1	9.19	39. <b>4</b> 0	3.38	15.0 18	.7 21		64.3	1 .8.1	~ o.	380	υ. Σ	7 10.0		
ND NOT DETERMINED	<u>م</u>																		

M
Š
2
FAR
[24

		TEST	VIT	WHT	PROTEIN 14% MB	EIN HB	FLR	WET	FLR	PEAK	MIX				HIX	Ä	COAF			
		Ē	200	9	E115	0 10	£λΩ	112	Hot	TIME		1 1		ABS	TIME	Š	7.	U		
VARIETY OR	хгр	<u>-</u>	7 P	2		1 T L	1 4 9	075	1100											
NIIMBER	BPA	LB/BU		SEC	جن	*	*	جن	مو	Z			CLASS	,,	Z					
ND 686	2	61.0	95	393			70.0		0.38	20.5		133	7.7AB	61.2						
ND 087	2	6.79	2	404			68.3		0.39	14.5			7.0	62.3						
NODE AND DO	2 2	9	. 0	432	14.2		65.0	38.7	0.39	22.2	26.0		8.0AB	67.0	2.35	10.01	1013 8	8.5	9.0 10.0	0 10.0
HORIZAN			) C	717			67.5		0.42	25.2	20.5		7.3AB	63.1						
ממא	2 5		5 6	0 0 0	10.0	12.7	61.5			7.5	26. B		7.7	64.1						
RUSS	3 :	0.10	r (	000	9 6	4.00					16.3		. 0	64.3	6					
SHARP	£	6 L . Y	7 6	716	r	7.7	- (				1 0		2.0	,						
BONJA	웊	58.9	9	391	13.5	17.8	68.3		0.40	0.47	27.5		. OAB	7.70	2 4					
STOA	QK	59.5	97	423	14.6	13.9	66.2		0.39	20.5	29.7		۵. ۵	63.0	CC.					
TORKTON	CH	60.5	95	415	13.9	13.2	67.5		0.38	24.2	23.2		7.7AB	63.3	85					
VERDE	G X	60.2	8	435	13.7	12.8	9.69		0.38	20.8	32.0		9.0	60.3	. 25					
2371	2	58.7	92	387	14.4	13.4	9.89		0.42	19.0	22.3		7.0	61.9	25					
2275	£	6.09	94	438	13.4	12.4	68.1		0.41	16.5	27.2	15	7.3	63.5						
2398	2	60.5	88	387	13.1	12.6	69.1	35.3	0.44	20.3	20.3	11	7.3AB	62.1	20			æ. æ.	.8 10.	
TOT	NED																			
	ILB 3-	· AC B	ATON	IA, I	RNEST	AND	HCNEA	AL GROWN	E-C	3 STA	STATIONS.	-					•			

#### EXHIBIT E - STATEMENT OF THE BASIS OF THE APPLICANT'S OWNERSHIP

Dr. Richard C. Frohberg, an employee of the North Dakota Agricultural Experiment Station and North Dakota State University, is a plant breeder who developed 'Ernest' the hard red spring wheat cultivar for which Plant Variety Protection is hereby sought. The employee by agreement and because of the condition of the use of facilities and funds of the North Dakota Agricultural Experiment Station and North Dakota State University has assigned all ownership rights to 'Ernest' hard red spring wheat to the North Dakota Agricultural Experiment Station and North Dakota State University.

North Dakota State University on behalf of the North Dakota Agricultural Experiment Station has assigned all ownership to the NDSU Research Foundation. The NDSU Research Foundation is a nonprofit corporation set up to own and manage the intellectual property of the North Dakota State University.